

SNIPPER

Woody Plant Deflowering Hormone

Contains Indole-3-butyric Acid Deflowering Hormonal Solution Applied Internally by TREE TECH Microinjection System for Promoting Premature Abscission of Male and/or Female Flowers.

ACTIVE INGREDIENT:

Indole-3-butyric acid	4.0%
INERT INGREDIENTS	96.0%
	<hr/>
	100.0%

NET CONTENTS: Each microinjection unit contains 0.2 grams a.i indole-3-butyric acid deflowering hormone.

U.S. Patent No. 3,051,723

SNIPPER is a Reg. TM of Florida Silvics, Inc.
Morrison, FL 32668

EPA Reg. No. 64014-13
EPA Est. No. 64014-FL-001

STOP READ THE LABEL BEFORE USE

FOR USE BY COMMERCIAL ARBORISTS
(APPLICATORS)

KEEP OUT OF REACH OF CHILDREN.

CAUTION

SEE REAR PANEL FOR STATEMENTS OF
PRACTICAL TREATMENT AND OTHER
PRECAUTIONARY STATEMENTS

NET CONTENTS: 5 mL per microinjection unit
100 microinjection units
(500 mL total) per case

Sold by:
FLORIDA SILVICS, INC.
(dba Tree Tech Microinjection Systems)
950 S.E. 215th Avenue
Morrison, FL 32668 USA

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING : Causes moderate eye injury. Avoid contact with skin, eyes or clothing. Harmful if inhaled or absorbed through the skin. Remove contaminated clothing and wash before reuse. Wash thoroughly with soap and water after handling, before eating, drinking and using tobacco.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are listed below.

USER SAFETY RECOMMENDATIONS

Users should :

1. Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
2. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
3. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

STATEMENT OF PRACTICAL TREATMENT

In case of ingestion call a physician or poison control center. Have the patient lie down and keep quiet.

IF IN EYES : Flush eyes with plenty of water. Call a physician if irritation persists.

IF ON SKIN : Remove contaminated clothing and immediately wash affected area with soap and water. If irritation occurs, get medical attention.

IF INHALED : Remove to fresh air. Get medical attention.

IF SWALLOWED : Call a physician or poison control center immediately. Induce vomiting by giving the patient 1 or 2 glasses of water and touching the back of the throat with finger. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person.

ENVIRONMENTAL HAZARDS

Do not apply directly to any body of water. Do not contaminate water by cleaning of equipment or disposal of wastes. Use for tree microinjection only as a pre-bloom or in-bloom application.

AVISO :

PRECAUCION AL USUARIO : Si Usted no puede leer o entender ingles, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

(TO THE USER : if you cannot read or understand English, do not use this product until the label has been fully explained to you.)

PHYSICAL/CHEMICAL HAZARDS

Do not use or store near heat or open flame.

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING . For any requirements specific to your State or Tribe, consult the agency responsible.

AGRICULTURAL USE REQUIREMENTS

Use this product in accordance with its labeling and with the Worker Protection Standard, 40 CFR 170. This standard contains requirements for the protection of agricultural workers on farms, forest nurseries, and greenhouses, and handlers or agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this labeling about personal protective equipment and restricted entry intervals.

The requirements in this box only apply to the uses of this product that are covered by the Workers Protection Standard (WPS).

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

THE RESTRICTED-ENTRY INTERVAL (REI)
FOR THIS PRODUCT IS "0" HOURS .

SNIPPER deflowering hormonal solution is for use by commercial applicators on sweet gum trees in commercial and residential landscapes, interior and exterior plantscapes, roadsides, gardens, parks, golf courses, lawns or grounds and other areas where sweet gum trees are grown. SNIPPER deflowering hormonal solution will be translocated throughout the vascular system of the plant.

GENERAL DIRECTIONS

Applying SNIPPER using the Tree Tech microinjection system :

SNIPPER label and microinjection instructions must be read and understood prior to use or installation of Tree Tech Microinjection Systems microinjection units. Failure to follow these directions may lead to injury to the installer or other persons as well as mechanical or phytotoxic damage to treated trees. The following instructions must be heeded to ensure proper and effective use of the microinjection units containing SNIPPER deflowering hormonal solution.

1. Do not inject trees that are less than two inches in diameter.
2. Do not inject trees within two weeks of any other spray or soil chemical treatment.
3. Do not treat trees that are suffering from stress such as lack of moisture or herbicide damage.
4. This product is not to be used on trees which will produce food within the year following treatment.
5. Protective eyewear and rubber or neoprene gloves must be worn while handling or installing the microinjection unit to prevent accidental contact with the eyes or skin.
6. When properly installed and activated, the microinjection unit generates internal pressure resulting in the flow of SNIPPER solution through the dispenser tube. The microinjection unit must never be activated unless installed correctly and securely in the tree to be treated.
7. Microinjection units containing SNIPPER may require up to several minutes or more to empty depending on the health of the treated tree and local weather conditions. Never assume that microinjection units have depressurized completely because they appear nearly empty or empty. When removing microinjection units, individuals must wear proper eye protection and rubber or neoprene gloves. The individual should then cover the microinjection unit with one hand near the point of insertion into the stem while grasping the barrel end of the microinjection unit with the other hand. The microinjection unit should be turned slightly as it is slowly withdrawn from the tree.
8. After microinjection units are removed from treated trees they must be discarded into the heavy-duty plastic disposal bag included in each case of microinjection units. The bag should be properly sealed and placed in the original carton. Sealed cartons should be returned freight prepaid to Tree Tech Microinjection Systems, 950 S.E. 215th Ave., Morrison, FL 32668 for disposal.

Installing Microinjection Units :

1. Determine the number of microinjection units to be installed based upon the recommended dosage rate as administered by proper circumferential spacing of microinjection units at 4-inch intervals around the stem. Unless otherwise noted, microinjection units should be installed in the stem and root flares near the ground line, i.e., 2.0-to-8.0 in. or 5-to-20 cm, from the soil surface.



SHERRILLtree®
TREE GEAR. TREE PEOPLE.

800-525-8873

SherrillTree.com

2. Using a cordless electric drill (600-to-800 rpm capacity is preferred) with a sharp, clean 11/64 in. (0.4 cm) bit, the installer should drill a hole at the correct stem circumference spacing to a depth of 3/8-to-1/2 in. (1.0-to-1.3 cm) into the wood (xylem) under the bark. A slight downwardly angle is recommended for more complete drainage of the microinjection unit
3. After reaching the proper depth, the drill bit should be withdrawn carefully to avoid dislodging bark fragments around the exterior opening of the hole. Insert the microinjection unit into the hole. Placing the plastic installation cap over the rear barrel end, strike the cap with a plastic hammer to seat the microinjection unit firmly into the hole. If the microinjection unit is not properly positioned in the hole, strike the cap again until correctly seated. By striking the microinjection unit, the frontal dispenser tip is forced into the funnel-shaped section dislodging an internal septum which allows the SNIPPER solution to flow from the microinjection unit into the tree. When the microinjection unit is positioned correctly in the tree and the septum is dislodged, remove the cap. Push the rear barrel portion of the unit downwardly until the locking mechanism is engaged. This pressurizes the microinjection unit and assists in the movement of SNIPPER into the vascular system of the tree.
4. Each hole should be drilled and a microinjection unit installed without delay. After the unit is properly seated, it should be activated. This sequence minimizes the flow of tree sap or resin into the hole prior to SNIPPER microinjection.
5. When properly installed and activated, the microinjection unit generates internal pressure resulting in the flow of SNIPPER solution through the dispenser tube. The microinjection unit must never be activated unless installed correctly and securely in the tree to be treated.
6. Microinjection units containing SNIPPER may require up to several minutes or more to empty depending on the health of the treated tree and local weather conditions, Microinjection units should be removed upon evacuation of SNIPPER deflowering hormonal solution.

APPLICATION: Apply in all cases when plant root systems are established and actively growing. Applications should be applied at recommended intervals to minimize seed formation.

Tree	Target	Microinjection unit Spacing Interval Around Stem Circumference	Time of Injection
Ash Black locust Black walnut Cherry Cottonwood Crabapple Gingko Hackberry Hickory Honey locust Maple Oak Olive Persimmon Plum Tree of Heaven Sweet gum Sycamore	Flowers (male and/or female)	One microinjection unit every 4 inches	After flower buds break and as flowers approach maximum size

GENERAL INFORMATION

SNIPPER deflowering hormonal solution is for use by commercial applicators on sweet gum trees. Not for use on sweet gum trees for research purposes.

CONDITIONS OF SALE

THE DIRECTIONS ON THIS LABEL WERE DETERMINED THROUGH RESEARCH TO BE APPROPRIATE FOR THE CORRECT USE OF THIS PRODUCT. THIS PRODUCT HAS BEEN TESTED UNDER DIFFERENT ENVIRONMENTAL CONDITIONS BOTH INDOORS AND OUTDOORS UNDER CONDITIONS SIMILAR TO THOSE THAT ARE ORDINARY AND CUSTOMARY WHERE THE PRODUCT IS TO BE USED. INSUFFICIENT CONTROL OF PESTS OR PLANT INJURY MAY RESULT FROM THE OCCURRENCE OF EXTRAORDINARY OR UNUSUAL CONDITIONS, OR FROM FAILURE TO FOLLOW LABEL DIRECTIONS. IN ADDITION, FAILURE TO FOLLOW LABEL DIRECTIONS MAY CAUSE INJURY TO ANIMALS, MAN, AND DAMAGE TO THE ENVIRONMENT. TREE TECH MICROINJECTION SYSTEMS OFFERS, AND THE BUYER ACCEPTS AND USES, THIS PRODUCT SUBJECT TO THE CONDITIONS THAT EXTRAORDINARY OR UNUSUAL ENVIRONMENTAL CONDITIONS, OR FAILURE TO FOLLOW LABEL DIRECTIONS ARE BEYOND THE CONTROL OF TREE TECH MICROINJECTION SYSTEMS, AND ARE THEREFORE, THE RESPONSIBILITY OF THE BUYER.

1. Tree Tech Microinjection Systems warrants that this product conforms to the chemical description on the label and is reasonably fit for the use under average conditions when used strictly in accordance with the directions on the labeling. Tree Tech Microinjection Systems does not make or authorize any agent or representative to make any other warranty, guarantee or representation, express or implied, concerning this product. Specifically, NO IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR ANY PARTICULAR PURPOSE IS MADE.

2. Critical and unforeseeable factors beyond the control of Tree Tech Microinjection Systems prevent it from eliminating all risks in connection with the use of this product. Such risks include, but are not limited to, damage to plants to which the product is applied, lack of complete control over the handling and application of this product, and damage caused by movement to other plants or crops. Such risks occur even though the product is reasonably fit under average conditions for the uses stated on the labeling and even though label directions are followed. Buyer and user acknowledge and assume all risks and liability (except those assumed by Tree Tech Microinjection Systems under 1 above) resulting from handling, storage and use of this product.

3. Precautions stated on the labeling should be followed to avoid hazardous exposure to the product. Neither Tree Tech Microinjection Systems nor its employees or distributors will be liable for any damages resulting from improper use of the microinjection units

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal of microinjection units.

Do not apply this product through any type of irrigation system.

Open dumping is prohibited.

Store microinjection units at room temperature (45 degrees F -to- 74 degrees F). Do not freeze. Do not contaminate water, food, or feed by storage or disposal.

STORAGE: Store product in the closed, original container in a cool, dry, locked place out of the reach of children.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site according to local regulations or at an approved waste disposal facility.

MICROINJECTION UNIT CONTAINER DISPOSAL
Do not re-use microinjection units. Used microinjection units should be placed in the heavy-duty plastic bag which accompanies each case of microinjection units. The bag should be properly sealed, placed into the original shipping carton and returned freight prepaid for disposal to Tree Tech Microinjection Systems, 950 S.E. 215th Ave., Morriston, FL 32696.

SNIPPER is a registered trademark of Florida Silvics, Inc.

Label revised 10/2002



SHERRILLtree®
TREE GEAR. TREE PEOPLE.

800-525-8873

SherrillTree.com